Site Assessment Report – 2013 – YPS-052

Site Data

Site	YPS-052
Sample Date	July 15 2013
Latitude	N 64.01888
Longitude	W 139.1275
Altitude	1312 ft
Feature Name	Hunker Creek
Stream Order	4

Site Photograph



Upstream view

Context Map

Refer to Klondike River Watershed Aquatic Health Monitoring Sites 2013 map.

Assignment of the Test Site to a Group

The test site is assigned to a benthic insect community group (Group) based on the 2013 Yukon CABIN model (BEAST Prediction Results). The site is assigned to the Group for which it has the highest probability of belonging based on habitat attributes.

Predictor	Altitude (ft), Longitude, LC-Bryoids (%), LC-Broadleaf open (%), LC-Mixed wood open					
Variables	(%), LC-Wetland herbaceous (%), Precipitation Feb (mm), Precipitation March (mm),					
	Precipitation June (mm), Precipitation July (mm), Rainfall June (mm), April Max temp					
	(C), Average Depth (cm), Average Velocity (m/s)					
Predicted Group						
	2					
Group		1	2	3	4	5
Probability that the site belongs 15.0%		39.6%	9.8%	34.3%	1.2%	
to each Group		13.070	33.070	5.070	34.370	1.2/0

Habitat Attributes of Site

This table reports on how the habitat predictor variables measured at the test site compare to the mean habitat predictor variables for the reference sites in the same group.

Variable	Site	Reference Mean	Reference SD
Longitude	-139.128	-136.93	2.75
Altitude (ft)	1312	2134.49	899.68
Depth-Avg (cm)	30.6	31.44	19.67
Velocity-Avg (m/s)	0.4	0.43	0.26
Precip. FEB (mm)	32.8	28.51	7.47
Precip. MAR (mm)	31.3	26.48	7.73
Precip. JUN (mm)	54.3	57.14	13.59
Precip. JUL (mm)	67.5	73.01	17.74
Rainfall JUN (mm)	51.9	49.32	11.37
April Max Temp (C)	-3.7	0.93	4.20
Broadleaf Open (%)	0.00	0.14	0.34
Bryoids (%)	0.47	0.31	0.61
Mixedwood Open (%)	0.23	0.75	1.44
Wetland Herb (%)	0.00	0.11	0.31

Summary Results of the Benthic Invertebrate Data

The table presents the summary values of the benthic community of the test site compared to expected values and the average benthic community of the reference sites.

	Test Site	Reference Average	Reference Standard Deviation	Expected Families (RIVPACS)
Total Abundance	79	265.34	160.63	
Total No. of Taxa	13	11.52	4.32	11.7

This table shows how the benthic community of the test site compares to the average of the reference sites to which it is being compared. Both the presence of certain families and their abundance (the number of individuals found) are compared.

Family (bold = characteristic of group)	RIVPACS Prediction (probability)	Test Site (# of individuals)	Reference Average	Reference Standard Deviation
Chironomidae	0.99	18	98	68
Baetidae	0.8	16	27	35
Simuliidae	0.76	0	16	31
Nemouridae	0.74	0	17	24
Heptageniidae	0.64	2	19	27
Sperchontidae	0.53	0	3	7
Tipulidae	0.51	1	2	3


Using the predictive model with the RIVAPCS weighted probability of the sites predicted membership, we would expect 11.7 families at this site, and 13 families were observed.

Of the seven families expected to occur (P>0.5), three were absent. Of the six families characteristic of the group to which this site was predicted, three were absent (Simuliidae, Nemouridae and Chloroperlidae). Overall, abundance was below the normal range, primarily due to low numbers of Chironomidae, while richness was slightly above expected.

Visual Reports on Site Condition

This figure displays three site assessment graphs which show the site relative to the group of reference sites to which it is compared. The three axes (or vectors) represent three dimensional space, and the probability bands (from center) are 75 - 90% (Similar to Reference), 90 - 99% (Mildly Divergent from Reference), 99 - 99.9% (Divergent from Reference), and outside 99.9% (Highly Divergent from Reference).

A 0-75% probability band was provided in past annual monitoring reports. Programming is currently being developed in order to incorporate this function into the CABIN model, however at this time the 0-75% band cannot be included here.



Assessment of Overall Site Condition

Vector 1 vs Vector 2	Similar to Reference
Vector 1 vs Vector 3	Mildly Divergent from Reference Condition
Vector 2 vs Vector 3	Mildly Divergent from Reference Condition
Overall	Mildly Divergent from Reference Condition

Using CABIN, YPS-052 is assessed as Mildly Divergent from Reference Condition.

Field Measurements

The following table shows general information collected at the site as well as water quality parameters when available. All measurements reflect conditions at the time the site visit was conducted.

Variable		
Site	YPS-052	
Sample Year	2013	
Status	Test	
Bankfull Width (m)	8.5	
Wetted Width (m)	5.6	
Channel Depth Average (cm)	30.6	
Channel Depth Max (cm)	42	
Slope (m/m)	0.0075	
Max Water Velocity (m/s)	0.48	
Average Water Velocity (m/s)	0.36	
Substrate Embeddedness	1/2 embedded	
Dominant Substrate-1st	6.4 – 12.8 cm (cobble)	
Dominant Substrate-2nd	0.2 – 1.6 cm (gravel)	
Surrounding Substrate Material	<0.1 cm (silt)	
Pools	Present	
Rapids	Absent	
Riffles	Present	
Straight Run	Present	
Canopy Coverage (%)	1-25	
Periphyton Coverage		
Macrophyte Coverage (%)	None	
Riparian Vegetation-Coniferous	Absent	
Riparian Vegetation-Deciduous	Present	
Riparian Vegetation-Grasses/Ferns	Present	
Riparian Vegetation-Shrubs	Present	
General Conductivity (μS/cm)	627	
Specific Conductance (μS/cm)	780	
DO (mg/L)	9.65	
рН (рН)	7.6	
TDS (mg/L)		
TSS (mg/L)		
Air Temp (Degrees Celsius)	28	
Water Temp (Degrees Celsius)	14.74	
Turbidity (NTU)		