# Site Assessment Report – 2013 – YPS- 431

## Site Data

Site	YPS- 431
Sample Date	Aug 8, 2013
Latitude	N 63.1912
Longitude	W 138.8425
Altitude	1227 ft
Feature Name	Scroggie Creek
Stream Order	6

# Site Photograph



Downstream view

# **Context Map**

Refer to Stewart River Watershed Aquatic Health Monitoring Sites 2013 map.

## Assignment of the Test Site to a Group

The test site is assigned to a benthic insect community group (Group) based on the 2013 Yukon CABIN model (BEAST Prediction Results). The site is assigned to the Group for which it has the highest probability of belonging based on habitat attributes.

Predictor	Altitude (ft), Longitude, LC-Bryoids (%), LC-Broadleaf open (%), LC-Mixed wood open (%), LC-					
Variables	Wetland herbaceous (%), Precipitation Feb (mm), Precipitation March (mm), Precipitation  June (mm), Precipitation July (mm), Rainfall June (mm), April Max temp (C), Average Depth					
	(cm), Average Velocity (m/s)					
Predicted Group	4					
Group		1	2	3	4	5
Probability that the each Group	site belongs to	18.7%	27.2%	11.8%	41.2%	1.0%

#### **Habitat Attributes of Site**

This table reports on how the habitat predictor variables measured at the test site compare to the mean habitat predictor variables for the reference sites in the same group.

Variable	Site	Reference Mean	Reference SD
Longitude	-138.843	-137.45	2.65
Altitude (ft)	1227	2296.81	838.01
Depth-Avg (cm)	31.0	29.80	14.62
Velocity-Avg (m/s)	0.7	0.52	0.32
Precip. FEB (mm)	36.4	29.34	11.79
Precip. MAR (mm)	35.7	27.46	11.91
Precip. JUN (mm)	60.9	53.49	18.49
Precip. JUL (mm)	76.9	65.85	22.37
Rainfall JUN (mm)	57.0	48.44	16.06
April Max Temp (C)	-2.6	-0.98	3.38
Broadleaf Open (%)	0.13	0.38	1.31
Bryoids (%)	0.64	0.54	1.04
Mixedwood Open (%)	1.73	0.77	2.87
Wetland Herb (%)	0.02	0.14	0.46

#### **Summary Results of the Benthic Invertebrate Data**

The table presents the summary values of the benthic community of the test site compared to expected values and the average benthic community of the reference sites.

	Test Site	Reference Average	Reference Standard Deviation	Expected Families (RIVPACS)
Total Abundance	238	2059.44	1572.86	
Total No. of Taxa	11	12.95	4.37	11.8

#### **Detailed Results of the Benthic Data**

This table shows how the benthic community of the test site compares to the average of the reference sites to which it is being compared. Both the presence of certain families and their abundance (the number of individuals found) are compared.


Family (bold = characteristic of group)	RIVPACS Prediction (probability)	Test Site (# of individuals)	Reference Average	Reference Standard Deviation
Chironomidae	0.98	31	911	827
Baetidae	0.78	99	233	309
Nemouridae	0.74	0	130	184
Simuliidae	0.75	3	207	367
Heptageniidae	0.64	5	120	216
Sperchontidae	0.54	0	35	60
Tipulidae	0.52	0	19	41

Using the predictive model with the RIVAPCS weighted probability of the sites predicted membership, we would expect 11.8 families at this site, and 11 families were observed. Of the seven families expected to occur (P>0.5), three (Nemouridae, Sperchontidae, Tipulidae), were absent. Of the five families characteristic of the group to which this site was predicted only Nemouridae were absent. The site is dominated by Baetidae rather than Chironomidae. Abundance was lower than the normal range, but richness was similar to what was expected.

#### **Visual Reports on Site Condition**

This figure displays three site assessment graphs which show the site relative to the group of reference sites to which it is compared. The three axes (or vectors) represent three dimensional space, and the probability bands (from center) are 75 - 90% (Similar to Reference), 90 - 99% (Mildly Divergent from Reference), 99 - 99.9% (Divergent from Reference), and outside 99.9% (Highly Divergent from Reference).

A 0-75% probability band was provided in past annual monitoring reports. Programming is currently being developed in order to incorporate this function into the CABIN model, however at this time the 0-75% band cannot be included here.



#### **Assessment of Overall Site Condition**

Vector 1 vs Vector 2	Mildly Divergent from Reference Condition
Vector 1 vs Vector 3	Mildly Divergent from Reference Condition
Vector 2 vs Vector 3	Similar to Reference
Overall	Mildly Divergent from Reference Condition

Using CABIN, YPS-431 is assessed as Mildly Divergent from Reference Condition.

## **Field Measurements**

The following table shows general information collected at the site as well as water quality parameters when available. All measurements reflect conditions at the time the site visit was conducted.

Variable	
Site	YPS-431
Sample Year	2013
Status	Test
Bankfull Width (m)	14.2
Wetted Width (m)	11
Channel Depth Average (cm)	31
Channel Depth Max (cm)	60
Slope (m/m)	0.01
Max Water Velocity (m/s)	1.33
Average Water Velocity (m/s)	0.69
Substrate Embeddedness	1/4 embedded
Dominant Substrate-1st	6.4 – 12.8 cm (cobble)
Dominant Substrate-2nd	12.8 – 25.6 cm (cobble)
Surrounding Substrate Material	0.1 – 0.2 cm (sand)
Pools	Present
Rapids	Absent
Riffles	Present
Straight Run	Present
Canopy Coverage (%)	1-25
Periphyton Coverage	
Macrophyte Coverage (%)	None
Riparian Vegetation-Coniferous	Present
Riparian Vegetation-Deciduous	Present
Riparian Vegetation-Grasses/Ferns	Present
Riparian Vegetation-Shrubs	Present
General Conductivity (μS/cm)	157
Specific Conductance (μS/cm)	216
DO (mg/L)	10.73
рН (рН)	7.9
TDS (mg/L)	
TSS (mg/L)	
Air Temp (Degrees Celsius)	17
Water Temp (Degrees Celsius)	10.65
Turbidity (NTU)	