# Site Assessment Report – 2013 – YPS- 544

# Site Data

Site	YPS- 544
Sample Date	July 15, 2013
Latitude	N 64.0294
Longitude	W 139.1784
Altitude	1177 ft
Feature Name	Hunker Creek
Stream Order	4

# Site Photograph



Upstream view

# **Context Map**

Refer to Klondike River Watershed Aquatic Health Monitoring Sites 2013 map.

### Assignment of the Test Site to a Group

The test site is assigned to a benthic insect community group (Group) based on the 2013 Yukon CABIN model (BEAST Prediction Results). The site is assigned to the Group for which it has the highest probability of belonging based on habitat attributes.

Predictor	Altitude (ft), Longitude, LC-Bryoids (%), LC-Broadleaf open (%), LC-Mixed wood open (%), LC-					
Variables	Wetland herbaceous (%), Precipitation Feb (mm), Precipitation March (mm), Precipitation June (mm), Precipitation July (mm), Rainfall June (mm), April Max temp (C), Average Depth  (cm), Average Velocity (m/s)					
Predicted Group	2					
Group		1	2	3	4	5
Probability that the site belongs to each Group		12.6%	38.8%	10.0%	37.1%	1.3%

#### **Habitat Attributes of Site**

This table reports on how the habitat predictor variables measured at the test site compare to the mean habitat predictor variables for the reference sites in the same group.

Variable	Site	Reference Mean	Reference SD
Longitude	-139.178	-136.93	2.75
Altitude (ft)	1177	2134.49	899.68
Depth-Avg (cm)	19.0	31.44	19.67
Velocity-Avg (m/s)	0.4	0.43	0.26
Precip. FEB (mm)	32.5	28.51	7.47
Precip. MAR (mm)	31.0	26.48	7.73
Precip. JUN (mm)	54.0	57.14	13.59
Precip. JUL (mm)	67.1	73.01	17.74
Rainfall JUN (mm)	51.5	49.32	11.37
April Max Temp (C)	-3.6	0.93	4.20
Broadleaf Open (%)	0.00	0.14	0.34
Bryoids (%)	0.45	0.31	0.61
Mixedwood Open (%)	0.23	0.75	1.44
Wetland Herb (%)	0.00	0.11	0.31

#### **Summary Results of the Benthic Invertebrate Data**

The table presents the summary values of the benthic community of the test site compared to expected values and the average benthic community of the reference sites.

	Test Site	Reference Average	Reference Standard Deviation	Expected Families (RIVPACS)
Total Abundance	40	265.34	160.63	
Total No. of Taxa	6	11.52	4.32	11.9

#### **Detailed Results of the Benthic Data**

This table shows how the benthic community of the test site compares to the average of the reference sites to which it is being compared. Both the presence of certain families and their abundance (the number of individuals found) are compared.


Family (bold = characteristic of group)	RIVPACS Prediction (probability)	Test Site (# of individuals)	Reference Average	Reference Standard Deviation
Chironomidae	0.99	12	98	68
Baetidae	0.81	18	27	35
Simuliidae	0.77	0	16	31
Nemouridae	0.76	0	17	24
Heptageniidae	0.66	0	19	27
Sperchontidae	0.54	0	3	7
Tipulidae	0.52	0	2	3
Empididae	0.51	0	4	10

Using the predictive model with the RIVAPCS weighted probability of the sites predicted membership, we would expect 11.9 families at this site, but 6 families were observed. Of the eight families expected to occur (P>0.5), only two were present (Chironomidae and Baetidae). Of the five families characteristic of the group to which this site was predicted three were absent. Total abundance was low compared to the normal range, and richness was lower than expected.

#### **Visual Reports on Site Condition**

This figure displays three site assessment graphs which show the site relative to the group of reference sites to which it is compared. The three axes (or vectors) represent three dimensional space, and the probability bands (from center) are 75 - 90% (Similar to Reference), 90 - 99% (Mildly Divergent from Reference), 99 - 99.9% (Divergent from Reference), and outside 99.9% (Highly Divergent from Reference).

A 0-75% probability band was provided in past annual monitoring reports. Programming is currently being developed in order to incorporate this function into the CABIN model, however at this time the 0-75% band cannot be included here.



## **Assessment of Overall Site Condition**

Vector 1 vs Vector 2	Similar to Reference
Vector 1 vs Vector 3	Mildly Divergent from Reference Condition
Vector 2 vs Vector 3	Mildly Divergent from Reference Condition
Overall	Mildly Divergent from Reference Condition

Using CABIN, YPS-544 is assessed as Mildly Divergent from Reference Condition.

## **Field Measurements**

The following table shows general information collected at the site as well as water quality parameters when available. All measurements reflect conditions at the time the site visit was conducted.

Variable	
Site	YPS-544
Sample Year	2013
Status	Test
Bankfull Width (m)	8.9
Wetted Width (m)	5.8
Channel Depth Average (cm)	19
Channel Depth Max (cm)	31
Slope (m/m)	0.005
Max Water Velocity (m/s)	0.63
Average Water Velocity (m/s)	0.42
Substrate Embeddedness	1/2 embedded
Dominant Substrate-1st	<0.1 cm (silt)
Dominant Substrate-2nd	3.2 – 6.4 cm (pebble)
Surrounding Substrate Material	<0.1 cm (silt)
Pools	Absent
Rapids	Absent
Riffles	Present
Straight Run	Present
Canopy Coverage (%)	1-25
Periphyton Coverage	
Macrophyte Coverage (%)	None
Riparian Vegetation-Coniferous	Absent
Riparian Vegetation-Deciduous	Present
Riparian Vegetation-Grasses/Ferns	Present
Riparian Vegetation-Shrubs	Present
General Conductivity (μS/cm)	527
Specific Conductance (μS/cm)	648
DO (mg/L)	10.22
рН (рН)	7.8
TDS (mg/L)	
TSS (mg/L)	
Air Temp (Degrees Celsius)	25
Water Temp (Degrees Celsius)	15.17
Turbidity (NTU)	