# Site Assessment Report – 2013 – YPS- 565

## Site Data

Site	YPS- 565
Sample Date	Aug 1, 2013
Latitude	N 61.5254
Longitude	W 139.3159
Altitude	2414 ft
Feature Name	Quill Creek
Stream Order	4

## Site Photograph



**Downstream View** 

## **Context Map**

Refer to White River Watershed Aquatic Health Monitoring Sites 2013 map.

## Assignment of the Test Site to a Group

The test site is assigned to a benthic insect community group (Group) based on the 2013 Yukon CABIN model (BEAST Prediction Results). The site is assigned to the Group for which it has the highest probability of belonging based on habitat attributes.

Predictor	Altitude (ft), Longitude, LC-Bryoids (%), LC-Broadleaf open (%), LC-Mixed wood open (%), LC-					
Variables	Wetland herbaceous (%), Precipitation Feb (mm), Precipitation March (mm), Precipitation June (mm), Precipitation July (mm), Rainfall June (mm), April Max temp (C), Average Depth  (cm), Average Velocity (m/s)					
Predicted Group	3					
Group		1	2	3	4	5
Probability that the each Group	site belongs to	28.4%	8.7%	34.0%	27.6%	1.3%

#### **Habitat Attributes of Site**

This table reports on how the habitat predictor variables measured at the test site compare to the mean habitat predictor variables for the reference sites in the same group.

Variable	Site	Reference Mean	Reference SD
Longitude	-139.316	-135.66	3.18
Altitude (ft)	2414	2756.11	719.61
Depth-Avg (cm)	38.0	32.11	15.81
Velocity-Avg (m/s)	1.5	0.58	0.29
Precip. FEB (mm)	36.8	36.14	23.93
Precip. MAR (mm)	33.8	33.13	21.04
Precip. JUN (mm)	58.8	64.67	18.69
Precip. JUL (mm)	76.0	78.30	20.81
Rainfall JUN (mm)	56.2	52.72	13.46
April Max Temp (C)	3.1	1.38	3.74
Broadleaf Open (%)	0.22	0.68	1.62
Bryoids (%)	0.72	0.37	0.84
Mixedwood Open (%)	0.00	0.96	1.72
Wetland Herb (%)	0.00	0.03	0.10

### **Summary Results of the Benthic Invertebrate Data**

The table presents the summary values of the benthic community of the test site compared to expected values and the average benthic community of the reference sites.

	Test Site	Reference Average	Reference Standard Deviation	Expected Families (RIVPACS)
Total Abundance	4	567.00	737.13	
Total No. of Taxa	2	10.56	6.13	11.8

This table shows how the benthic community of the test site compares to the average of the reference sites to which it is being compared. Both the presence of certain families and their abundance (the number of individuals found) are compared.


Family (bold = characteristic of group)	RIVPACS Prediction (probability)	Test Site (# of individuals)	Reference Average	Reference Standard Deviation
Chironomidae	0.98	2	162	142
Nemouridae	0.75	0	175	168
Simuliidae	0.72	2	37	55
Baetidae	0.71	0	92	109
Heptageniidae	0.65	0	311	288
Sperchontidae	0.53	0	13	22
Chloroperlidae	0.50	0	32	38
Empididae	0.50	0	16	29
Tipulidae	0.50	0	5	10

Using the predictive model with the RIVAPCS weighted probability of the sites predicted membership, we would expect 11.8 families at this site, but only 2 families were observed. Of the nine families expected to occur (P>0.5), only two (Chironomidae and Simuliidae) were present. Only two of the six families characteristic of the group to which this site was predicted were present and with very low abundance (represented by only two organisms each). Total abundance was well below the normal range and richness was much lower than expected.

#### **Visual Reports on Site Condition**

This figure displays three site assessment graphs which show the site relative to the group of reference sites to which it is compared. The three axes (or vectors) represent three dimensional space, and the probability bands (from center) are 75 - 90% (Similar to Reference), 90 - 99% (Mildly Divergent from Reference), 99 - 99.9% (Divergent from Reference), and outside 99.9% (Highly Divergent from Reference).

A 0-75% probability band was provided in past annual monitoring reports. Programming is currently being developed in order to incorporate this function into the CABIN model, however at this time the 0-75% band cannot be included here.



## **Assessment of Overall Site Condition**

Vector 1 vs Vector 2	Highly Divergent from Reference Condition
Vector 1 vs Vector 3	Analysis Not Required
Vector 2 vs Vector 3	Analysis Not Required
Overall	Highly Divergent from Reference Condition

Using CABIN, YPS-565 is assessed as Highly Divergent from Reference Condition. Stream flow was very high during sampling and this likely effected stream velocity measurements as well as invertebrate sampling. Model results for this site are not considered representative at this time. In addition the site location may not be representative of the stream reach and therefore this site will not be re-sampled.

## **Field Measurements**

The following table shows general information collected at the site as well as water quality parameters when available. All measurements reflect conditions at the time the site visit was conducted.

Variable	
Site	YPS-565
Sample Year	2013
Status	Test
Bankfull Width (m)	90
Wetted Width (m)	4
Channel Depth Average (cm)	38
Channel Depth Max (cm)	52
Slope (m/m)	0.02
Max Water Velocity (m/s)	2.5
Average Water Velocity (m/s)	1.53
Substrate Embeddedness	1/4 embedded
Dominant Substrate-1st	6.4 – 12.8 cm (cobble)
Dominant Substrate-2nd	0.2 – 1.6 cm (gravel)
Surrounding Substrate Material	<0.1 cm (silt)
Pools	Present
Rapids	Present
Riffles	Absent
Straight Run	Absent
Canopy Coverage (%)	1-25
Periphyton Coverage	
Macrophyte Coverage (%)	None
Riparian Vegetation-Coniferous	Present
Riparian Vegetation-Deciduous	Present
Riparian Vegetation-Grasses/Ferns	Present
Riparian Vegetation-Shrubs	Present
General Conductivity (μS/cm)	249
Specific Conductance (μS/cm)	369
DO (mg/L)	11.53
рН (рН)	7.9
TDS (mg/L)	
TSS (mg/L)	
Air Temp (Degrees Celsius)	18
Water Temp (Degrees Celsius)	8.04
Turbidity (NTU)	