Site Assessment Report 2014 – YPS-431

Site Data

Site	YPS-431
Sample Date	Jul 23, 2014
Latitude	63.19125 N
Longitude	138.84254 W
Altitude	1227
Feature Name	Scroggie Creek
Stream Order	6

Site Photograph – Aerial View



Context Map

Refer to Stewart River Aquatic Health Monitoring Sites 2014 map.

Site Sampling History

This site has been sampled on two previous occasions, in 2010 and 2013.

Assignment of the Test Site to a Group

The test site is assigned to a benthic insect community group (Group) based on the 2013 Yukon CABIN model (BEAST Prediction Results). The site is assigned to the Group for which it has the highest probability of belonging based on habitat attributes.

Predictor Variables	Altitude (ft), Longitude, LC-Bryoids (%), LC-Broadleaf open (%), LC-Mixed wood open (%), LC-Wetland herbaceous (%), Precipitation Feb (mm), Precipitation March (mm), Precipitation June (mm), Precipitation July (mm), Rainfall June (mm), April Max temp (C),					
	Average Depth (cm), Average Velocity (m/s)					
Predicted Group	4					
Group		1	2	3	4	5
Probability that the seach Gro	Ŭ	24.6%	27.9%	11.2%	35.6%	0.7%

Habitat Attributes of Site

This table reports on how the habitat predictor variables measured at the test site compare to the mean habitat predictor variables for the reference sites in the same group.

Variable	Site	Reference Mean	Reference SD
Longitude	-138.84	-137.45	2.65
Altitude (ft)	1227	2296.81	838.01
Depth-Avg (cm)	44.00	29.80	14.62
Velocity-Avg (m/s)	0.68	0.52	0.32
Precip. FEB (mm)	36.39	29.34	11.79
Precip. MAR (mm)	35.74	27.46	11.91
Precip. JUN (mm)	60.86	53.49	18.49
Precip. JUL (mm)	76.91	65.85	22.37
Rainfall JUN (mm)	57.02	48.44	16.06
April Max Temp (C)	-2.61	-0.98	3.38
Broadleaf Open (%)	0.13	0.38	1.31
Bryoids (%)	0.64	0.54	1.04
Mixedwood Open (%)	1.73	0.77	2.87
Wetland Herb (%)	0.02	0.14	0.46

Summary Results of the Benthic Invertebrate Data

The table presents the summary values of the benthic community of the test site compared to expected values and the average benthic community of the reference sites.

	Test Site	Reference Average	Reference Standard Deviation	Expected Families (RIVPACS)
Total Abundance	401.00	2059.44	1572.86	
Total No. of Taxa	11.00	12.95	4.37	11.3

This table shows how the benthic community of the test site compares to the average of the reference sites to which it is being compared. Both the presence of certain families and their abundance (the number of individuals found) are compared.

Family (bold = characteristic of group)	RIVPACS Prediction (probability)	Test Site (# of individuals)	Reference Average	Reference Standard Deviation
Chironomidae	0.98	135	911	827
Baetidae	0.74	97	233	309
Simuliidae	0.72	6	207	367
Nemouridae	0.71	0	130	184
Heptageniidae	0.60	2	120	216
Sperchontidae	0.51	0	35	60
Tipulidae	0.50	0	19	41

Using the predictive model with the RIVAPCS weighted probability of the sites predicted membership, we would expect 11.3 families at this site, but 11 families were observed.

Total abundance is well below average for this group and outside the normal range. Richness (total number of families) is as expected.


Of the 7 families expected to occur (P>0.5), 3 (Nemouridae, Sperchontidae, and Tipulidae) were absent.

Of the five families characteristic of the group to which this site was predicted, one was absent (Nemouridae).

Visual Reports on Site Condition

This figure displays three site assessment graphs which show the site relative to the group of reference sites to which it is compared. The three axes (or vectors) represent three dimensional space, and the probability bands (from center) are 75 - 90% (Similar to Reference), 90 - 99% (Mildly Divergent from Reference), 99 - 99.9% (Divergent from Reference), and outside 99.9% (Highly Divergent from Reference).

A 0-75% probability band was provided in past annual monitoring reports. Programming is currently being developed in order to incorporate this function into the CABIN model, however at this time the 0-75% band cannot be included here.



Assessment of Overall Site Condition

Vector 1 vs Vector 2	Mildly Divergent from Reference Condition
Vector 1 vs Vector 3	Mildly Divergent from Reference Condition
Vector 2 vs Vector 3	Similar to Reference
Overall	Mildly Divergent from Reference Condition

Using CABIN, YPS-431 is assessed as Mildly Divergent from Reference Condition.

Field Measurements

The following table shows general information collected at the site as well as water quality parameters when available. All measurements reflect conditions at the time the site visit was conducted.

Site	YPS-431
Sample Year	2014
Status	Test
Bankfull Width (m)	12.8
Wetted Width (m)	11.9
Channel Depth Average (cm)	44
Channel Depth Max (cm)	51
Slope m/m	0.005
Slope (%)	0.5
Max Water Velocity (m/s)	1
Average Water Velocity (m/s)	0.68
Dominant Substrate-1st	12.8 – 25.6 cm (cobble)
Dominant Substrate-2nd	6.4 – 12.8 cm (cobble)
Surrounding Substrate Material	<0.1 cm (silt)
Substrate Embeddedness	1/4 Embedded
Pools	Present
Rapids	Absent
Riffles	Present
Straight Run	Present
Canopy Coverage (%)	1-25
Periphyton Coverage	
Macrophyte Coverage (%)	None
Riparian Vegetation-Coniferous	Present
Riparian Vegetation-Deciduous	Present
Riparian Vegetation-Grasses/Ferns	Present
Riparian Vegetation-Shrubs	Present
General Conductivity (μS/cm)	143.9
Specific Conductance (μS/cm)	201.8
DO (mg/L)	9.03
рН (рН)	7.9
TDS (mg/L)	115
TSS (mg/L)	81
Air Temp (Degrees Celsius)	17
Water Temp (Degrees Celsius)	10.6
Turbidity (NTU)	54